

BLACK SPRUCE

***Picea mariana* (Miller) B.S.P.**

Plant Symbol = PIMA

Contributed by: USDA NRCS National Plant Data Center



© Kenneth J. Sytsma
University of Wisconsin – Madison
Wisconsin State Herbarium

Alternate Names

Bog spruce, swamp spruce, shortleaf black spruce

Uses

The primary use of black spruce wood is for pulp. Lumber is of secondary importance because of the relatively small size of the trees. The trees and wood also are used for fuel, Christmas trees, and other products (beverages, medical salves, aromatic distillations). Black spruce is the provincial tree of Newfoundland.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Description

General: Trees to 25 meters tall (often shrub-like near tree-line), the crown narrowly conic to spire-like or "irregularly subcylindric;" branches short and drooping, frequently layering; twigs not pendent, slender, yellow-brown, hairy; bark gray-brown. Needles evergreen, 0.6-1.5(-2) cm long, 4-angled, stiff and blunt-tipped, waxy and pale blue-green. Seed cones 1.5-2.5(-3.5) cm long, fusiform, purple-brown at maturity; cone scales fan-shaped, broadest near apex, 8-12 mm long, rigid, margin at apex irregularly toothed. Native. The common name refers to the dark (blackish) foliage.

Variation within the species: Black spruce shows north-south clinal variation in photoperiodism, productivity, and other traits, and discrete variants are not recognized at present, except for one. Fernald (1950) recognized "depressed" and "trailing" alpine forms at the rank of *forma*, but Roland and Smith (1969) and Rouleau and Lamoureux (1992) have treated them as var. *semiprostrata*. These occur in Keewatin, Quebec, Newfoundland, and Nova Scotia:

- *Picea mariana* var. *mariana*
- *Picea mariana* var. *semiprostrata* (Peck) Teeri
- Synonym: *Picea mariana* forma *semiprostrata* (Peck) Blake

However, these are of little significance to NRCS and conservation personnel in the United States.

Black spruce hybridizes with red spruce (*Picea rubens*) on disturbed sites in eastern Canada and with white spruce in a few places. Reports of natural hybrids between black spruce and white spruce apparently remain unverified.

Distribution: Black spruce occurs across the northern range of North America, from Alaska, Yukon, and British Columbia eastward to Nova Scotia, Newfoundland, New Brunswick, and Québec. It also occurs in the northeastern United States and sporadically in Minnesota, Wisconsin, and Michigan. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

Adaptation

In muskegs, bogs, bottomlands, and relatively dry peatlands; at 0-1500 meters. Black spruce usually grows on wet organic soils but productive stands also grow over deep humus, clays, loams, sands, coarse

till, and shallow soil mantles. It is often a postfire pioneer on both uplands and peatlands. In fire-prone areas, such as upland ridges, fire usually results in the immediate reestablishment and eventual dominance of black spruce, because it produces seed at an early age.

Establishment

Black spruce may produce a few cones as early as 10 years of age, but maximum production occurs between 100 and 200 years. Heavy seed years occur at intervals of 2-6 years and peak crops every 4 years over most of the range. Fires open the cones and accelerate seed release for periods of 60 days to 2-3 years, depending on fire intensity. Viable seed may be dispersed from a cone for up to 25 years, providing a seed supply after fires occur.

Sphagnum mosses can provide a continuously moist seedbed for black spruce, but growth of seedling growth may be slow here because of inadequate nutrients. Feathermosses may provide a suitable seedbed during wet years but may dry out before penetration by the seedling root. Moist mineral soils usually provide good seedbeds, but exposed mineral soil may be too waterlogged or subject to frost heaving in some low-lying areas. Fires that completely remove the surface organic layer usually provide good seedbeds. Seedling mortality seems to be highest on burned duff and lowest on moss and mineral soil surfaces with adequate moisture. Seedlings develop in as little as 10% of full light intensity, but survival and growth are much better in the open.

Layering is an important means of reproduction in black spruce on some sites, especially where rapidly growing mosses cover lower branches. Such layering develops most abundantly in the more open-grown stands and less frequently in dense stands with higher wood volume. It also is common at tree line, probably as a result of depression of the lower branches by snow, and accounts for the presence of "candelabrum" spruce, a circular clump originating from one individual with the tallest tree in the center.

The average maximum age for black spruce is about 200 years, but ages up to 300 years have been reported.

Management

Black spruce grows more slowly than many associated trees and shrubs, and mature trees in spruce-fir stands apparently respond better to release than white spruce and subalpine fir. Many

intermediate and suppressed black spruce in swamp stands, however, die after heavy cutting

Clearcutting in strips or patches is generally considered to be the best silvicultural system for managing black spruce. Satisfactory reestablishment after clearcutting requires an adequate seed source and often some kind of site preparation. Uneven-aged or all-aged management is best applied on poor sites where stands are windfirm and have abundant layering.

Eastern dwarf-mistletoe causes serious problems for black spruce in the Lake States and eastern Canada. The spruce budworm and various other insects are damaging. Black spruce is easily killed by both ground and crown fires. Peatland stands have a low risk except during very dry periods

Cultivars, Improved and Selected Materials (and area of origin)

Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

- Fernald, M.L. 1950. *Gray's manual of botany*. Ed. 8. American Book Co., New York.
- Little, E.L., Jr. and S.S. Pauley. 1958. A natural hybrid between black and white spruce in Minnesota. *Amer. Midl. Naturalist* 60:202-211.
- Morgenstern, E.K. and J.L. Farrar. 1964. *Introgressive hybridization in red spruce and black spruce*. Univ. Toronto, Fac. Forest., Techn. Rep. 4.
- Roland, A.E. and E.C. Smith. 1969. *The flora of Nova Scotia*. Nova Scotia Museum, Halifax.
- Rouleau, E. and G. Lamoureux 1992. *Atlas of the vascular plants of the island of Newfoundland and the islands of St. Pierre et Miquelon*. Groupe Fleurbec, St. Henri de Levis, Quebec.
- Sytsma, K.J. 2004. *Wisconsin State Herbarium*. (<http://www.botany.wisc.edu/wisflora/scripts/detail.asp?SpCode=PICMAR>, 20 December 2004). University of Wisconsin, Madison.
- Taylor, R.J. 1993. *Picea*. Pp. 369-373 in: *Flora of North America, north of Mexico*. vol. 2. Oxford Univ. Press, New York.

Teeri, J.A. 1969. The phytogeography of subalpine black spruce in New England. *Rhodora* 71:1-6.

Viereck, L.A. and W.F. Johnston 1990. *Picea mariana*. Pp. 227-237 in: R.M. Burns and B.H. Honkala Silvics of North America. vol 1. USDA Forest Service Agric. Handbook 654, Washington, D.C.

Prepared By:

Guy Nesom

Formerly BONAP, North Carolina Botanical Garden
University of North Carolina, Chapel Hill, North Carolina

Species Coordinator:

Gerald Guala

USDA NRCS National Plant Data Center, Baton Rouge, Louisiana

Edited: 20Dec2004 rln; 060802 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's [TARGET Center](#) at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about [Civil Rights at the Natural Resources Conservation Service](#).